

# ERGA Assembly Report

v24.10.15

Tags: Other[INVALID TAG]

TxID	7917
ToLID	<b>fAtrSpal_hap1</b>
Species	Alligator gar
Class	Actinopteri
Order	Semionotiformes

Genome Traits	Expected	Observed
Haploid size (bp)	509,031,378	1,168,890,252
Haploid Number	34 (source: ancestor)	28
Ploidy	2 (source: ancestor)	2
Sample Sex	Unknown	Unknown

## EBP metrics summary and curation notes

Obtained EBP quality metric for collapsed: 7.7.Q61

The following metrics were automatically flagged as below EBP recommended standards or different from expected:

- . Observed Haploid size (bp) has >20% difference with Expected
- . Observed Haploid Number is different from Expected

### Curator notes

- . Interventions/Gb: 31
- . Contamination notes: ""
- . Other observations: "The assembly of Alligator gar (fAtrSpal) has been generated as part of the Vertebrate Genomes Project (<https://vertebrategenomesproject.org/>). Both haplotypes were analyzed and manually improved using Pretext. "

## Quality metrics table


Metrics	Pre-curation collapsed	Curated collapsed
Total bp	1,123,788,851	1,168,890,252
GC %	40.47	40.62
Gaps/Gbp	57.84	53.9
Total gap bp	13,000	12,600
Scaffolds	345	334
Scaffold N50	55,314,901	55,314,901
Scaffold L50	9	9
Scaffold L90	28	26
Contigs	410	397
Contig N50	24,563,650	25,314,049
Contig L50	16	16
Contig L90	57	50
QV	61.7709	61.3836
Kmer compl.	96.4638	97.0348
BUSCO sing.	88.0%	96.7%
BUSCO dupl.	0.6%	0.6%
BUSCO frag.	3.5%	0.6%
BUSCO miss.	7.9%	2.1%

Warning! BUSCO versions or lineage datasets are not the same across results:

BUSCO: 5.8.2 (euk\_genome\_met, metaeuk) / Lineage: actinopterygii\_odb12 (genomes:75, BUSCOs:7207)


BUSCO: 6.0.0 (euk\_genome\_min, miniprot) / Lineage: actinopterygii\_odb12 (genomes:75, BUSCOs:7207)

# HiC contact map of curated assembly




collapsed [\[LINK\]](#)

# K-mer spectra of curated assembly




Distribution of k-mer counts per copy numbers found in asm





Distribution of k-mer counts coloured by their presence in reads/assemblies

# Post-curation contamination screening



TAPAs summary Graph



**collapsed.** Bubble plot circles are scaled by sequence length, positioned by coverage and GC proportion, and coloured by taxonomy. Histograms show total assembly length distribution on each axis.

# Data profile

Data	Long reads	Arima
Coverage	NA	NA

# Assembly pipeline

- **Hifiasm**
  - |\_ *ver*: 0.19.5-r593
  - |\_ *key param*: NA
- **purge\_dups**
  - |\_ *ver*: 1.2.5
  - |\_ *key param*: NA
- **YaHS**
  - |\_ *ver*: 1.2
  - |\_ *key param*: NA

# Curation pipeline

- **PretextMap**
  - |\_ *ver*: 0.1.9
  - |\_ *key param*: NA
- **PretextView**
  - |\_ *ver*: 0.2.5
  - |\_ *key param*: NA

Submitter: Benjamin Istace

Affiliation: Genoscope

Date and time: 2025-11-05 18:14:56 CET